

Tropical linear mappings on the plane

M.J. de la Puente^{*†}

July 16, 2009

Abstract

In this paper we fully describe all *tropical linear mappings in the tropical projective plane* $\mathbb{TP}^2$, that is, maps from the tropical plane to itself given by tropical multiplication by a 3×3 matrix A with entries in $\mathbb{T}$. First we will allow only real entries in the matrix A and, only at the end of the paper, we will allow some of the entries of A equal $-\infty$. The mapping f_A is continuous and piecewise-linear in the classical sense. In some particular cases, the mapping f_A is a parallel projection onto the set spanned by the columns of A . In the general case, after a change of coordinates, the mapping collapses at most three regions of the plane onto certain segments, called antennas, and is a parallel projection elsewhere (theorem 3).

In order to study f_A , we may assume that A is normal, i.e., $I \leq A \leq 0$, up to changes of coordinates. A given matrix A admits infinitely many normalizations. Our approach is to define and compute a unique normalization for A (which we call *canonical normalization*) (theorem 1) and then always work with it, due both to its algebraic simplicity and its geometrical meaning.

On $\mathbb{R}^n$, any $n \in \mathbb{N}$, some aspects of tropical lineal maps have been studied in [5]. We work in $\mathbb{TP}^2$, adding a geometric view and doing everything explicitly. We give precise pictures.

Inspiration for this paper comes from [3, 5, 7, 11, 24]. We have tried to make it self-contained. Our preparatory results present noticeable relationships between the algebraic properties of a given matrix A (normal idempotent matrix, permutation matrix, etc.) and classical geometric properties of the points spanned by the columns of A (classical convexity and others); see theorem 2 and corollary 1. As a by-product, we compute all the tropical square roots of normal matrices of a certain type; see corollary 3. This is, perhaps, a curious result in tropical algebra. Our final aim is, however, to give a precise description of the mapping $f_A : \mathbb{TP}^2 \rightarrow \mathbb{TP}^2$. This is particularly easy when two tropical triangles arising from A (denoted $\mathcal{T}_A$ and $\mathcal{T}^A$) fit as much as possible. Then the action of f_A is easily described on each cell of the cell decomposition $\mathcal{C}^A$; see theorem 3.

Normal matrices play a crucial role in this paper. The tropical powers of normal matrices of size $n \in \mathbb{N}$ satisfy $A^{\odot n-1} = A^{\odot n} = A^{\odot n+1} = \dots$. This statement can be traced back, at least, to [24], and appears later many times, such

^{*}Departamento de Algebra, Facultad de Matemáticas, Universidad Complutense, 28040-Madrid, Spain

[†]Partially supported by UCM research group 910444.

as [1, 2, 5, 8, 9]. In lemma 1, we give a direct proof of this fact, for $n = 3$. But now the equality $A^{\odot 2} = A^{\odot 3}$ means that the columns of $A^{\odot 2}$ are three fixed points of f_A and, in fact, *any point spanned by the columns of $A^{\odot 2}$ is fixed by f_A* . Among 3×3 normal matrices, the *idempotent* ones (i.e., those satisfying $A = A^{\odot 2}$) are particularly nice: we prove that the columns of such a matrix tropically span a set which is classically compact, connected and *convex* (lemma 2 and corollary 1). In our terminology, it is a *good tropical triangle*.

1 Introduction, Notations and Background on Tropical Mathematics

Many results on finite dimensional tropical linear algebra (spectral theory, etc.) have been published over the last 40 years and more; they are summarized in [1, 9, 13], where a wide bibliography can also be found. In this paper we will use the adjective *classical* as opposed to *tropical*. Most definitions in tropical mathematics just mimic the classical ones. However, tropical geometry is a peculiar one. Say an inhabitant of the tropical plane is disoriented. He/she takes a look at a compass and tries to spot the tropical cardinal points. There are only three: *east, north and south-west*! Accordingly, he/she will set the positive part of the three coordinate axes in the given directions, when doing geometry on the plane. He/she will find out that a generic tropical line in the tropical plane looks like a tripod (they have a vertex!), although some particular tropical lines look just like classical lines, see figure 1.




Figure 1: Tropical line with vertex at the point $(-2, -2)$.

If we happen to go down-town in a city designed by a tropical geometer, we will

find out that the shape of most blocs is that of a classical hexagon, with parallel opposite sides of slopes 0, 1, ∞ , see figure 2.




Figure 2: Downtown blocs in a tropical city.

The shortest path between two given points is made up of, at most, two classical segments with slopes 0, 1, ∞ . Moreover, the distance between the given points is the sum of the *integer lengths* (also called lattice length) of these segments. For instance, the integer length between the points $(-2, -2)$ and $(0, 0)$ is 2 (not $2\sqrt{2}$!) and the integer length between the points $(-5, -2)$ and $(0, 0)$ is $3 + 2 = 5$. This is indeed, a sort of Manhattan distance.

So, plane tropical geometry is a funny looking piecewise-linear geometry. And, by the way, why is it called tropical? Well, the explanation appears in [12, 14], etc. and we must add that some other names have also been used (for this or akin mathematics): max-plus, dioids, path algebra, extremal algebra, idempotent mathematics, etc.

Consider the set $\mathbb{R} \cup \{-\infty\}$ endowed with tropical addition $\oplus$ and tropical multiplication $\odot$, where these operations are defined as follows:

$$a \oplus b = \max\{a, b\}, \quad a \odot b = a + b,$$

for $a, b \in \mathbb{R} \cup \{-\infty\}$. Here, $-\infty$ is the neutral element for tropical addition and 0 is the neutral element for tropical multiplication. Notice that $a \oplus a = a$, for all a , i.e., tropical addition is *idempotent*. Notice also that a has no inverse with respect to $\oplus$.

We will work with $\mathbb{R} \cup \{-\infty\}$, which will be denoted $\mathbb{T}$ and will be called the *tropical semi-field*. We will write $\oplus$ or $\max$, (resp. $\odot$ or $+$) at our convenience.

In classical mathematics, we have a choice in geometry: affine or projective. The *tropical affine plane* is $\mathbb{T}^2$, where addition and multiplication are defined coordinate-wise. In the space $\mathbb{T}^3 \setminus \{(-\infty, -\infty, -\infty)\}$ we define an equivalence relation $\sim$ by letting $(p_1, p_2, p_3) \sim (q_1, q_2, q_3)$ if there exists $\lambda \in \mathbb{R}$ such that

$$\lambda \odot (p_1, p_2, p_3) = (\lambda + p_1, \lambda + p_2, \lambda + p_3) = (q_1, q_2, q_3).$$

The equivalence class of (p_1, p_2, p_3) is denoted $[p_1, p_2, p_3]$. The *tropical projective plane* is the set, $\mathbb{TP}^2$, of such equivalence classes. Notice that, at least, one of the coordinates of any point in $\mathbb{TP}^2$ must be finite.

We endow the tropical plane (either affine or projective) with the topology induced by the *Euclidean topology*. Thus, topological notions of a set S such as closure $\overline{S}$ will refer to this topology. It can be easily proved that $\mathbb{TP}^2$ is compact. In p. 9 below, we also define a *tropical norm* in the projective tropical plane. This norm gives rise to the Euclidean topology.

$\mathbb{TP}^2$ is a *compactification* of $\mathbb{T}^2$. Indeed, we have an injective mapping $\varphi : \mathbb{T}^2 \rightarrow \mathbb{TP}^2$ given by $(x, y) \mapsto [x, y, 0]$. The image of φ is open and dense. Now, for any $p = [x, y, z]$, we have $\varphi^{-1}(p) = (x - z, y - z)$, whenever $z \neq -\infty$. Taking $(x - z, y - z, 0)$ as a representative of p will be expressed by saying that *we work in $Z = 0$* . In other words, it is just a way of passing from the projective to the affine tropical plane. $\mathbb{TP}^2$ has three *boundary components*: these are the sets of points $[x, y, z]$ in $\mathbb{TP}^2$ where either $x = -\infty$ or $y = -\infty$ or $z = -\infty$.

The simplest objects in the tropical plane are lines. Given a tropical linear form

$$p_1 \odot X \oplus p_2 \odot Y \oplus p_3 \odot Z = \max\{p_1 + X, p_2 + Y, p_3 + Z\}$$

a *tropical line* consists of the points $[x, y, z] \in \mathbb{TP}^2$ where *the maximum is attained, at least, twice*, (this is the tropical analog of the classical vanishing point set). Denote this line by L_p , where $p = [p_1, p_2, p_3] \in \mathbb{TP}^2$.

Most lines in the tropical plane look like *tripods*. Indeed, if two coefficients are equal to $-\infty$, then L_p is a boundary component of $\mathbb{TP}^2$. If $p_j = -\infty$ for just one j then, in $Z = 0$, L_p is nothing but a classical slope-one line. If all p_j are real, the L_p is the union of three rays. The directions of these rays are West, South and North-East (just opposite to the cardinal directions of the tropical plane!) and these rays are emanating from the point $-p$, called the *vertex* of L_p . The latter is the generic case.

Let two points p, q in the tropical plane be given. The *tropical stable join* of p, q is defined as the limit, as ϵ tends to zero, of the tropical lines going through perturbed points $p^{v_\epsilon}, q^{v_\epsilon}$. Here, p^{v_ϵ} denotes a translation of p by a length- ϵ vector v_ϵ , see [12, 21]. We denote this line by pq .

Now, given two tropical lines L_p, L_q in the plane, the *stable intersection* of L_p, L_q , denoted $L_p \cap_{\text{st}} L_q$, is defined as the limit point, as ϵ tends to zero, of the intersection of perturbed lines $L_p^{v_\epsilon}, L_q^{v_\epsilon}$. Here, $L_p^{v_\epsilon}$ denotes a translation of L_p by a length- ϵ vector v_ϵ .

There exists a *duality* between lines and points since

$$q \in L_p \iff p \in L_q,$$

meaning that *the maximum $\max\{p_1 + q_1, p_2 + q_2, p_3 + q_3\}$ is attained, at least, twice*. This duality transforms stable join into stable intersection and conversely, i.e.,

$$L_p \cap_{\text{st}} L_q = r \iff pq = L_r,$$

for p, q, r in $\mathbb{TP}^2$.

The *tropical version of Cramer's rule* (see [21]) goes as follows: the stable intersection of the lines L_p and L_q is the point

$$[\max\{p_2 + q_3, q_2 + p_3\}, \max\{p_1 + q_3, q_1 + p_3\}, \max\{p_1 + q_2, q_1 + p_2\}].$$

Since the computation of this point is nothing but a *tropical version of the cross-product* of the triples p and q , we will denote it by $p \otimes q$ (this is not to be mixed up with $p \odot q = p + q$). Notice that $p \otimes q = q \otimes p$. In other words, the *tropical version of Cramer's rule* in the plane can be written as

$$L_p \cap_{\text{st}} L_q = p \otimes q \quad \text{and} \quad pq = L_{p \otimes q},$$

by duality. In particular, $-(p \otimes q)$ is the vertex of the line pq , a crucial fact that we use again and again.




Figure 3: Tropical line segments.

Given a subset U of points in $\mathbb{TP}^2$ (resp. $\mathbb{T}^2$), we can consider the *tropical span* of U , denoted $\text{span}(U)$, meaning the set of points $u \in \mathbb{TP}^2$ (resp. $\mathbb{T}^2$) which can be written as

$$u = \lambda_1 \odot u_1 \oplus \cdots \oplus \lambda_s \odot u_s = \max\{\lambda_1 + u_1, \dots, \lambda_s + u_s\},$$

for some $s \in \mathbb{N}$, $u_1, \dots, u_s \in U$, $\lambda_1, \dots, \lambda_s \in \mathbb{T}$, and not all λ_j equal to $-\infty$. The tropical span of two points p, q is called a *tropical segment*. We know that $\text{span}(p, q)$ is the union of the classical segments $\overline{p, -(p \otimes q)}$ and $\overline{-(p \otimes q), q}$. Similarly, the co-span of p, q is the union of the classical segments $\overline{p, (-p) \otimes (-q)}$ and $\overline{(-p) \otimes (-q), q}$.

The *tropical co-span* of U , denoted $\text{co-span}(U)$, is the set of points u which can be written as

$$u = \min\{\lambda_1 + u_1, \dots, \lambda_s + u_s\},$$

for some $s \in \mathbb{N}$, $u_1, \dots, u_s \in U$, $\lambda_1, \dots, \lambda_s \in \mathbb{R} \cup \{+\infty\}$, and not all λ_j equal to $+\infty$.

Given two points $p, q \in \mathbb{TP}^2$, we know that $-(p \otimes q)$ represents the vertex of the stable tropical line joining p and q . On the other hand, the stable intersection of the tropical lines with vertices at p, q is represented by the point $(-p) \otimes (-q)$. It turns out that the points $p, -(p \otimes q), q, (-p) \otimes (-q)$ are the vertices of a classical parallelogram, see figure 4.




Figure 4: Span and co-span of points p, q .

Another sort of duality is taking place here. Indeed, we may consider $\mathbb{R} \cup \{+\infty\}$ endowed with tropical addition $\ominus = \min$ and the same tropical multiplication $\odot$. The relationship between these two operations is $\max\{p, q\} = -\min\{-p, -q\}$, whence

$$p \oplus q = -(-p) \ominus (-q),$$

for $p, q \in \mathbb{R}$. This *max-min duality* appears in the literature, see [4, 8, 7], etc.

Why do we care about the co-span? A tropical triangle can be determined by three points, or by three lines. First, a tropical triangle $\mathcal{T}$ is determined by three points a, b, c . If the points are tropically collinear then $\mathcal{T}$ is not generic. We have

$$\mathcal{T} = \text{span}(a, b, c).$$

The *sides* of $\mathcal{T}$ are, by definition, the tropical lines ab, bc and ca . The vertices of the sides of $\mathcal{T}$ (as tropical lines) are $-(a \otimes b), -(b \otimes c)$ and $-(c \otimes a)$, again by the *tropical*

version of Cramer's rule. The properties of the triangle $\mathcal{T}$ depend on the configuration of the six points

$$a, b, c, -(a \otimes b), -(b \otimes c), -(c \otimes a), \quad (1)$$

which are all different, generically.

Three tropical lines L_p, L_q, L_r also determine a *tropical triangle*, $\mathcal{T}'$, which is generic if the lines are not tropically concurrent. We can write

$$\mathcal{T}' = \text{co-span}(-p, -q, -r).$$

The stable intersections (by pairs) of the lines L_p, L_q, L_r are called *vertices* of $\mathcal{T}'$. These points should not be mixed up with the vertices $-p, -q, -r$ of the lines. By the *tropical version of Cramer's rule*, the coordinates of the vertices of $\mathcal{T}'$ are $p \otimes q, q \otimes r$ and $r \otimes p$. The properties of $\mathcal{T}'$ depend on the configuration of the six points

$$p \otimes q, q \otimes r, r \otimes p, -p, -q, -r,$$

which are all different, in the generic case.

Tropical triangles are, in general, infinite unions of tropical segments. Indeed,

$$\mathcal{T} = \text{span}(a, b, c) = \bigcup_{s \in \text{span}(b, c)} \text{span}(a, s). \quad (2)$$

Therefore, tropical triangles are, in general, connected non-pure 2-dimensional sets. The non-generic case arises when the points a, b, c are tropically collinear. In addition, it is easy to check that tropical triangles are classically compact, both in $\mathbb{TP}^2$ and in $\mathbb{T}^2$.

It is not true, in general, that the stable intersection of the tropical lines ab and bc gives back the point b , and this makes *tropical triangles trickier than classical triangles*. For example, take $a = [3, 4, 6], b = [-2, 0, 8]$ and $c = [1, 1, 0]$. Then $a \otimes b = [12, 11, 3], b \otimes c = [9, 9, 1]$ and $ab \cap_{\text{st}} bc = [12, 13, 21] = [-1, 0, 8] \neq b$. The reader is encouraged to draw this example, in $Z = 0$.

This anomalous situation for tropical triangles has been studied in [3], where the definition of *good tropical triangle* has been given. Three points a, b, c define a *good tropical triangle* if, by stable join, they give rise to three tropical lines ab, bc, ca which, stably intersected by pairs, yield the original points a, b, c , i.e.,

$$ca \cap_{\text{st}} ab = a, \quad ab \cap_{\text{st}} bc = b, \quad bc \cap_{\text{st}} ca = c.$$

Good tropical triangles are characterized by six slack inequalities. Indeed, write the coordinates of (representatives of) a, b, c as the columns of a matrix $A = (a_{ij})$ so that c occupies the first column and a occupies the third. Write

$$\mathcal{T}_A = \text{span}(A). \quad (3)$$

Then theorem 2 in [3] tells us that $\mathcal{T}_A \subseteq \mathbb{TP}^2$ is a good tropical triangle if and only if

$$a_{12} - a_{22} \leq a_{13} - a_{23} \leq a_{11} - a_{21},$$

$$a_{23} - a_{33} \leq a_{21} - a_{31} \leq a_{22} - a_{32}, \quad (4)$$

$$a_{31} - a_{11} \leq a_{32} - a_{12} \leq a_{33} - a_{13}.$$

Write

$$A_0 = \begin{bmatrix} a_{11} - a_{31} & a_{12} - a_{32} & a_{13} - a_{33} \\ a_{21} - a_{31} & a_{22} - a_{32} & a_{23} - a_{33} \\ 0 & 0 & 0 \end{bmatrix}$$

in order to make drawings in $Z = 0$. It is easy to check that the six inequalities (4) imply the following *cardinal points condition in $Z = 0$* : $\text{col}(A_0, 1)$ represents the most eastwards point, $\text{col}(A_0, 2)$ represents the most northwards one, and $\text{col}(A_0, 3)$ represents the most south-westwards one, among the columns of A_0 . The converse is not true.




Figure 5: Good triangle determined by the matrix A .

In figure 5 we see the good tropical triangle determined by the matrix

$$A = A_0 = \begin{bmatrix} 0 & -5 & -7 \\ -3 & 0 & -6 \\ 0 & 0 & 0 \end{bmatrix}.$$

Actually, in $Z = 0$, good tropical triangles are nothing but classical hexagons, pentagons, quadrangles or triangles having slopes 0, 1 and ∞ , where the inequalities (4) provide the integer length of the sides. They are obtained by chopping off two corners, in a classical rectangle, see figure 6. In figure 7 we see a few good triangles for which some inequalities are equalities.




Figure 6: A good tropical triangle is a classical rectangle with two corners chopped off.




Figure 7: Some good tropical triangles.

In the classical plane $\mathbb{R}^2$ we have the following norm

$$\|p\| = \max\{\|p_1\|, \|p_2\|, \|p_1 - p_2\|\}, \quad p \in \mathbb{R}^2.$$

It is easy to check that $\|p\|$ is the integer length of the tropical segment $\text{span}(p, 0)$, if we identify $p = (p_1, p_2)$ with $[p_1, p_2, 0]$. For instance, $\|(-5, -2)\| = 5$, $\|(-3, 5)\| = 8$. The unit ball and some radii in it are shown in figure 8. Given real points $p, q \in \mathbb{R}^2$ the *tropical distance* between p and q is $\|p - q\|$, by definition. It is the integer length of the tropical segment $\text{span}(p, q)$. This is connected with the generalized Hilbert projective metric appearing in [7, 11, 15] and to the range seminorm of [10].




Figure 8: Axes and unit ball in the tropical the plane. Some rays are shown in dotted lines.

In $\mathbb{TP}^2$, let V be the tropical span of a finite family of points. The *projector map* (or *nearest point map*) $\rho : \mathbb{TP}^2 \rightarrow V$ satisfies $\rho \circ \rho = \rho$ and $\rho|_V = \text{id}_V$. For a point $p \in \mathbb{TP}^2 \setminus V$, the image $\rho(p)$ is computed as follows: fix a representative $p' \in \mathbb{T}^3$ of p and, for each generator v of V , choose a representative $v' \in \mathbb{T}^3$, optimal for the condition $v' \leq p'$ (meaning $v'_j \leq p'_j$, for $j = 1, 2, 3$ and equality is attained for, at least, one j). Tropically add all such v' s and then, take $\rho(p)$ to be the point in $\mathbb{TP}^2$ represented by the sum. In [7, 11, 15] it is shown that $\rho(p) \in V$ minimizes the tropical distance $\|p - q\|$, when q runs through V . In general, there are infinitely many points q in V minimizing such a distance, in addition to $\rho(p)$. Indeed, we consider tropical balls $B(d)$ centered at p of increasing radius d and take the minimum $d > 0$ such that the intersection $B(d) \cap V$ is non-empty. Then $B(d) \cap V$ is the set of minimizing points.

2 Matrices, mappings and pictures in $Z = 0$

All arrays will have entries in $\mathbb{T}$. Arrays will be denoted by capital letters A, B, C, N, P, Q , etc. Tropical matrix addition and multiplication are defined in the usual way, but using the tropical operations $\oplus$ and $\odot$, instead of the classical ones. Any array all whose entries are zero will be denoted by 0. Given two arrays of the same size $A = (a_{ij}), B = (b_{ij})$, we will write $A \leq B$ if $a_{ij} \leq b_{ij}$, for all i, j .

Our square matrices will have entries in $\mathbb{T}$, but *at least one entry in each row and in each column will be real*. If all the entries of a matrix A are real, we will say that A is a *real matrix*. We will deal with 3×3 matrices. The *tropical determinant* of a 3×3

matrix $A = (a_{ij})$ (also called *tropical permanent*) is defined as

$$|A|_{trop} = \max_{\sigma \in \Sigma_3} \{a_{1\sigma(1)} + a_{2\sigma(2)} + a_{3\sigma(3)}\},$$

where Σ_3 denotes the symmetric group in 3 symbols. A matrix is *tropically singular* if the maximum in the tropical determinant is attained, at least, twice. Otherwise the matrix is *tropically regular*, or it is said to have a *strong permanent*. These are all standard definitions.

Given a matrix A , the j -th column (resp. row) of A will be denoted $\text{col}(A, j)$ (resp. $\text{row}(A, j)$). The triple of diagonal entries of A will be denoted $\text{diag}(A)$. Moreover, if $t \in \mathbb{R}^3$, then $\text{diag}(t)$ will denote the matrix whose diagonal is t , the rest of entries being equal to $-\infty$; such matrices will be called *diagonal matrices*. A *permutation matrix* is a matrix obtained from a diagonal matrix, by permuting some of its rows or permuting some of its columns. A particular case is the *tropical identity matrix*, $I = \text{diag}(0)$. Another example is

$$P_{12} = \begin{bmatrix} -\infty & 0 & -\infty \\ 0 & -\infty & -\infty \\ -\infty & -\infty & 0 \end{bmatrix}.$$

Any permutation matrix P has a tropical inverse $P^{\odot -1}$, meaning $P \odot P^{\odot -1} = P^{\odot -1} \odot P = I$.

From now on, points in $\mathbb{TP}^2$ will be denoted by columns, for convenience. We often identify a 3×3 matrix A with the three points in $\mathbb{TP}^2$ represented by its columns.

The reader can easily check that left-multiplication by the matrix P_{12} exchanges coordinates X and Y :

$$P_{12} \odot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} y \\ x \\ z \end{bmatrix}.$$

A triple $t = (t_1, t_2, t_3) \in \mathbb{R}^3$ gives rise to a *translation* in $\mathbb{TP}^2$:

$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \mapsto \begin{bmatrix} X' \\ Y' \\ Z' \end{bmatrix} = \begin{bmatrix} t_1 + X \\ t_2 + Y \\ t_3 + Z \end{bmatrix} = \text{diag}(t) \odot \begin{bmatrix} X \\ Y \\ Z \end{bmatrix},$$

By a *change of projective coordinates* in the tropical projective plane $\mathbb{TP}^2$ we mean left-multiplying coordinates by a permutation matrix. Therefore, a change of projective coordinates amounts to the composite of a translation and a permutation of coordinates. Notice that right-multiplying A by a diagonal matrix does not change the columns of A in $\mathbb{TP}^2$; it only changes the representatives of them.

All pictures will be done in the affine tropical plane $Z = 0$. In order to do so, from a given matrix A we compute the matrix

$$A_0 = A \odot (\text{diag}(-\text{row}(A, 3))). \quad (5)$$

Conversely, if A_0 is a matrix having $\text{row}(A_0, 3) = 0$, then we recover A as follows:

$$A = A_0 \odot \text{diag}(-A_0). \quad (6)$$

Our aim is to describe the mapping $f_A : \mathbb{TP}^2 \rightarrow \mathbb{TP}^2$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto A \odot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \max\{a_{11} + x, a_{12} + y, a_{13} + z\} \\ \max\{a_{21} + x, a_{22} + y, a_{23} + z\} \\ \max\{a_{31} + x, a_{32} + y, a_{33} + z\} \end{bmatrix}.$$

First, notice that tropically proportional matrices A and $\lambda \odot A$ determine the same mapping $f_A = f_{\lambda \odot A}$, any $\lambda \in \mathbb{R}$. The simplest example of mapping f_A arises for $A = 0$, the mapping being constant. This is also true for $f_{A \otimes 0}$ and $f_{0 \otimes A}$.

The mapping f_A is obviously continuous and piecewise-linear. The image $\text{im } f_A$ is the tropical triangle spanned by A , meaning that it is spanned by the columns of A :

$$\mathcal{T}_A = \text{im } f_A = \text{span}(A). \quad (7)$$

If A is real, then *the mapping f_A is not surjective*, since no finite family of points with finite coordinates span the whole $\mathbb{TP}^2$; this is well-known (see, e.g., [23]). Moreover, if $r, s \in \mathbb{R}$ are negative and big enough, we have

$$A \odot \begin{bmatrix} 0 \\ r \\ s \end{bmatrix} = \text{col}(A, 1), \quad A \odot \begin{bmatrix} s \\ 0 \\ r \end{bmatrix} = \text{col}(A, 2), \quad A \odot \begin{bmatrix} r \\ s \\ 0 \end{bmatrix} = \text{col}(A, 3).$$

Therefore, f_A is *locally constant* on three big chunks of $\mathbb{TP}^2$, called *corners*. In particular, f_A is *not injective*.

Let us see how do these corners arise. First, the matrix A defines three tropical lines A_1, A_2, A_3 , because the j -th row of A provides a tropical linear form

$$a_{j1} \odot X \oplus a_{j2} \odot Y \oplus a_{j3} \odot Z = \max\{a_{j1} + X, a_{j2} + Y, a_{j3} + Z\}.$$

The vertices of A_1, A_2, A_3 are (represented by) the rows of $-A$, i.e., the columns of $-A^T$. Thus we have another tropical triangle here, namely

$$\mathcal{T}^A = \text{co-span}(-A^T). \quad (8)$$

Moreover, the lines A_1, A_2, A_3 (or, rather, the matrix A) induce a *cell decomposition* of $\mathbb{TP}^2$, denoted $\mathcal{C}^A$ (see [11] for an isomorphic cell decomposition). The decomposition $\mathcal{C}^A$ consists of, at most, 31 cells, and this is the generic case.

It has:

- ten 2-dimensional cells: one bounded cell, denoted B^A , the three already mentioned *corners* (denoted C_1^A, C_2^A, C_3^A), six unbounded cells (parallel to some tropical coordinate axis X, Y or Z),
- fifteen 1-dimensional cells: nine unbounded cells (parallel to some coordinate axis) and six bounded cells,
- six 0-dimensional cells or points.

Notice that the union of the bounded cells above is nothing but

$$\overline{B^A} = \mathcal{T}^A. \quad (9)$$

In figure 9 we find the 31 cells described above, and figure 10 represents the cell decomposition induced by the matrix

$$A = \begin{bmatrix} 0 & -1 & -5 \\ -4 & 0 & -2 \\ -1 & -4 & 0 \end{bmatrix}, \quad (-A^T)_0 = \begin{bmatrix} -5 & 2 & 1 \\ -4 & -2 & 4 \\ 0 & 0 & 0 \end{bmatrix}.$$




Figure 9: The 31 cells in the cell decomposition $\mathcal{C}^A$ induced by a matrix A .

The description of the mapping $f_A : \mathbb{TP}^2 \rightarrow \mathbb{TP}^2$ is particularly easy when the tropical triangles $\mathcal{T}_A$ and $\mathcal{T}^A$ fit as much as possible: then the action of f_A is easily described on each cell of the cell decomposition $\mathcal{C}^A$; see theorem 3.




Figure 10: Cell decomposition $\mathcal{C}^A$ induced by matrix A .

3 Normal matrices

By definition, a matrix A is *normal* if $\text{diag}(A) = 0$ and $A \leq 0$; in symbols,

$$I \leq A \leq 0 \quad (10)$$

see [5]. For any matrix A there exist permutation matrices P, Q such that the product

$$N = P \odot A \odot Q \quad (11)$$

is normal. The matrix N is called *a normalization of A* . The *Hungarian method* (see [5, 16, 20]) is an algorithm to obtain such N, P, Q . A matrix A admits several normalizations. Notice that the columns of A and the columns of $A \odot Q$ represent the same points in $\mathbb{TP}^2$, given perhaps in a different order. And the columns of N are a just a translation of those points.

As in classical mathematics, the product of matrices corresponds to the composite of mappings:

$$f_N = f_P \circ f_A \circ f_Q.$$

Now, f_P and f_Q are changes of projective coordinates, so that *in order to study the mapping f_A , we may assume that A is normal, up to changes of coordinates*.

A normal matrix A satisfies $I \leq A \leq 0$, and therefore

$$I \leq A \leq A^{\odot 2} \leq A^{\odot 3} \leq \dots \quad (12)$$

since tropical multiplication by any matrix is monotonic (because $\max$ and $+$ are monotonic). Moreover, for any natural m , $A^{\odot m+1} \leq A^{\odot m} \odot 0$ and the mapping $f_{A^{\odot m} \odot 0}$ is constant. In corollary 3 we see that the tropical powers of A are simpler than A (in the sense that they depend on fewer parameters), when A belongs to a particular class of normal matrices. This simplification carries over to the corresponding mappings

$$\text{id} = f_I, f_A, f_{A^{\odot 2}}, f_{A^{\odot 3}}, \dots, \text{const}.$$

In $\mathbb{TP}^2$, we consider the cell decomposition $\mathcal{C}^0$ induced by the zero matrix, i.e., the cell decomposition given by the tropical line L_0 . It has only three 2-dimensional cells (corners), which have the following description in $Z = 0$:

$$C_1^0 = \{0 < x, y < x\}, \quad C_2^0 = \{0 < y, x < y\}, \quad C_3^0 = \{x < 0, y < 0\}.$$

The *geometric meaning of normality* is the following: if A is a 3×3 normal matrix then,

$$\text{col}(A_0, j) \in \overline{C_j^0}, \quad \text{for all } j = 1, 2, 3. \quad (13)$$

Conversely, if A_0 is a 3×3 matrix with $\text{row}(A_0, 3) = 0$ satisfying (13), then A in (6) is normal.

Next we define several operators on matrices and then we study the relationship among them. Of course, we are particularly interested in these operators acting on normal matrices.

For any $k \in \mathbb{N}$, the tropical k -th power of A , denoted $A^{\odot k}$, takes normal matrices to normal matrices. The transpose A^T of a normal matrix A is a normal matrix. These operators commute with each other. Warning: $(-A)^{\odot 2} \neq A^{\odot 2}$, in general. Also, $(A_0)^{\odot 2} \neq (A^{\odot 2})_0$, in general.

We introduce the *tropical adjoint* of A , denoted $\hat{A}$. By definition, $\hat{A} = (\alpha_{ij})$, where α_{ij} is the tropical cofactor of a_{ji} . In other words,

$$\text{row}(\hat{A}, j) = \text{col}(A, j-1) \otimes \text{col}(A, j+1), \quad (14)$$

for $j = 1, 2, 3, \text{ mod } 3$. Last, we define an auxiliary matrix operator, $\check{A} = (\beta_{ij})$, by the formulas

$$\beta_{ii} = 0, \quad \beta_{ij} = a_{ik} + a_{kj},$$

if $i \neq j$ and $\{i, j, k\} = \{1, 2, 3\}$.

Lemma 1. *If A is 3×3 and normal, then*

1. $\check{A}$ is normal and $\hat{A} = A \oplus \check{A} = A^{\odot 2}$,
2. $\hat{A}$ is normal,
3. $A^{\odot 2} = A^{\odot 3}$,
4. the columns of $A^{\odot 2}$ represent fixed points of f_A ,
5. zero (the neutral element for tropical multiplication) is an eigenvalue of A .

Proof. A straightforward computation yields (1) and then (2) follows. Now, multiplication by A is a monotonic operator; so that the equality in (1) implies $A^{\odot 3} = \max\{A^{\odot 2}, A \odot \check{A}\}$. Now, a simple computation shows that $A \odot \check{A} = A \oplus \hat{A}$, whence $A^{\odot 3} = A^{\odot 2}$ follows. Finally, (4) follows from (3) and (5) follows from (4). $\square$

Lemma 1 follows from [24], where real matrices of any size n are considered. The so called *Kleene star of A* (or *strong closure of A*) is defined as

$$A^* = I \oplus A \oplus A^{\odot 2} \oplus A^{\odot 3} \oplus \dots,$$

if the limit exists, see[1, 6]. If A is a 3×3 normal matrix, then $A^* = A^{\odot 2}$, but we will not use this.

Lemma 2. *For a 3×3 normal matrix A , the following are equivalent:*

1. $\check{A} \leq A$,
2. $A = A^{\odot 2}$,
3. $\mathcal{T}_A$ is good.

Proof. The equivalence follows from lemma 1 and the six inequalities (4), letting $a_{jj} = 0$, for $j = 1, 2, 3$. Indeed, we obtain

$$\begin{aligned} a_{23} + a_{31} &\leq a_{21}, & a_{32} + a_{21} &\leq a_{31}, \\ a_{13} + a_{32} &\leq a_{12}, & a_{31} + a_{12} &\leq a_{32}, \\ a_{12} + a_{23} &\leq a_{13}, & a_{21} + a_{13} &\leq a_{23}. \end{aligned} \tag{15}$$

□

If $A = (a_{ij})$ is normal, we have

$$A_0 = \begin{bmatrix} -a_{31} & a_{12} - a_{32} & a_{13} \\ a_{21} - a_{31} & -a_{32} & a_{23} \\ 0 & 0 & 0 \end{bmatrix}.$$

By a translation, we can assume that $a_{13} = a_{23} = 0$, so that $\text{col}(A_0, 3) = 0$. Write

$$t_1 = -a_{31}, \quad t_2 = -a_{32}, \quad t_3 = a_{21} - a_{31}, \quad t_4 = a_{12} - a_{32}. \tag{16}$$

Then

$$t_1, t_2 \geq 0, \quad 0 \leq t_3, t_4 \leq \min\{t_1, t_2\} \tag{17}$$

and the t_j provide a *parameter space* for good tropical triangles, up to translation; see figure 6.

4 Canonical normalization

The geometric idea of canonical normalization is to center the figures at the origin of $Z = 0$. For each $d, d_1, d_2, d_3 \in \mathbb{R}$ with $d \geq 0$ and $-d \leq d_j$, for $j = 1, 2, 3$, the matrix

$$D(d, d_1, d_2, d_3) = \begin{bmatrix} 0 & -d - d_2 & -2d - d_3 \\ -2d - d_1 & 0 & -d - d_3 \\ -d - d_1 & -2d - d_2 & 0 \end{bmatrix} \tag{18}$$

is normal and it is easy to check that $D = D^{\odot 2}$. Notice the symmetric role played by d_1 with respect to X , d_2 with respect to Y and d_3 with respect to Z . We will use the matrices

$$D(d, d_1, d_2, d_3)_0 = \begin{bmatrix} d + d_1 & d & -2d - d_3 \\ -d & 2d + d_2 & -d - d_3 \\ 0 & 0 & 0 \end{bmatrix}, \quad (19)$$

$$(-D(d, d_1, d_2, d_3)^T)_0 = \begin{bmatrix} -2d - d_3 & d + d_1 - d_3 & d + d_1 \\ -d + d_2 - d_3 & -d - d_3 & 2d + d_2 \\ 0 & 0 & 0 \end{bmatrix}. \quad (20)$$




Figure 11: Tropical triangle given by the matrix $D(d, d_1, d_2, d_3)$, in $Z = 0$. Dotted lines are auxiliary.

Lemma 3 (Canonical normalization for an idempotent normal matrix). *If A is a 3×3 normal matrix such that $A = A^{\odot 2}$, then there exist unique $d, d_1, d_2, d_3 \geq 0$ and there exist unique permutation matrices P, Q such that $D(d, d_1, d_2, d_3) = P \odot A \odot Q$.*

Proof. By the geometric meaning of normality (13), a translation allows us to assume that $\text{col}(A_0, 3) = 0$. Then the triangle $\mathcal{T}_A$ is determined by the parameters $t_1, t_2, t_3, t_4 \geq 0$ as in (17).

$$A_0 = \begin{bmatrix} t_1 & t_4 & 0 \\ t_3 & t_2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad A = \begin{bmatrix} 0 & t_4 - t_2 & 0 \\ t_3 - t_1 & 0 & 0 \\ -t_1 & -t_2 & 0 \end{bmatrix}.$$

We take $d = \frac{1}{3}|t_3 - t_4|$, $d_3 = \min\{t_3, t_4\}$, $m = \max\{t_3, t_4\}$, $d_j = t_j - m$, for $j = 1, 2$ and $Q = \text{diag}(d_3 + 2d, d_3 + d, 0)$. Then $D(d, d_1, d_2, d_3) = Q^{\odot -1} \odot A \odot Q$. The uniqueness follows from the geometric meaning of the parameters d, d_1, d_2, d_3 . $\square$




Figure 12: Tropical triangle associated to a normal idempotent matrix.

Corollary 1. *A good tropical triangle is classically convex in $Z = 0$.*

Proof. Let $\mathcal{T}_A$ be a good tropical triangle, for some matrix A . By the paragraph after (11), a translation allows us to assume that A is normal. By lemma 3, we can assume that $A = D(d, d_1, d_2, d_3)$, for some $d, d_1, d_2, d_3 \geq 0$. To work in $Z = 0$, we consider the matrix A_0 shown above in (19). The columns of A_0 span $\mathcal{T}_A$, which is a classical hexagon, pentagon, rectangle or triangle of slopes 0, 1 and ∞ in $Z = 0$. By the tropical version of Cramer's rule, equality (14) and lemma 2, $\mathcal{T}_A$ it is actually the convex hull of the columns of A_0 and columns of $(-\hat{A}^T)_0 = -(A^{\odot 2})^T)_0 = (-A^T)_0$ shown in (20). $\square$

Lemma 4. *Given numbers $d, d_1, d_2, d_3 \in \mathbb{R}$, with $d \geq 0$ and $-d \leq d_j$, for $j = 1, 2, 3$, consider the normal matrix $A = D(d, d_1, d_2, d_3)$. The following are equivalent:*

1. $A \neq A^{\odot 2}$,
2. $d_j < 0$, for some $j = 1, 2, 3$,
3. in $Z = 0$, $\mathcal{T}_A$ is not classically convex.

Proof. To check for convexity in $Z = 0$, consider the matrix A_0 in expression (19). Say, $j = 1$. If $d_1 < 0$, then any point in the classical segment $a = \overline{p, q}$, with $p = [d + d_1, 2d + d_1 + d_2, 0]^T$ and $q = \text{col}(A_0, 2) = [d, 2d + d_2, 0]^T$, prevents $\mathcal{T}_A$ from being convex; see figure 13. $\square$

We will say that a is an *antenna* of $\mathcal{T}_A$.




Figure 13: Tropical triangle with antenna due to $d_1 < 0$.

In the hypothesis of the former lemma, $\mathcal{T}_A$ admits a *cell decomposition* having, at most, 13 cells, and this is the generic case:

- one 2-dimensional cell,
- six 1-dimensional cells,
- six 0-dimensional cells.

The closure of any 1-dimensional cell disrupting the convexity of $\mathcal{T}_A$ in $Z = 0$ is called an *antenna*, as in [7]. The union of points in the antennas of $\mathcal{T}_A$ will be denoted $\text{ant}(\mathcal{T}_A)$. Each j with $d_j < 0$ yields an antenna in $\mathcal{T}_A$. The integer length of this antenna is, at most, d .

Corollary 2. *Given $d, d_1, d_2, d_3 \in \mathbb{R}$, $d \geq 0$, $-d \leq d_j$, for $j = 1, 2, 3$, let $A = D(d, d_1, d_2, d_3)$. Then $\mathcal{T}_{A^{\odot 2}}$ is the maximal convex set contained in $\mathcal{T}_A$, in $Z = 0$.*

Proof. We know that A is normal. If $A = A^{\odot 2}$, we just have corollary 1. Otherwise $A^{\odot 2} = (A^{\odot 2})^{\odot 2}$, by lemma 1, so that $\mathcal{T}_{A^{\odot 2}}$ is convex in $Z = 0$, by lemma 2 and corollary 1.

Now we compute $A^{\odot 2} = (b_{ij})$ using (1) of lemma 1, obtaining $a_{ij} = b_{ij}$ unless $(i, j) = (1, 3), (2, 1)$ or $(3, 2)$. If, say, $d_1 < 0$, then

$$b_{32} = -2d - d_1 - d_2 > a_{32} = -2d - d_2.$$

Then, $\text{col}((A^{\odot 2})_0, 2) = [d + d_1, 2d + d_1 + d_2, 0]^T$ and $\text{col}(A_0, 1) = [d + d_1, -d, 0]^T$ so that both points belong to the same classical line $X = d + d_1$, meaning that the antenna in $\mathcal{T}_A$ caused by the inequality $d_1 < 0$ no longer appears in $\mathcal{T}_{A^{\odot 2}}$. $\square$

Suppose $d \geq 0$ and $-d \leq d_j$, all j . The former corollary tells us that *squaring the normal matrix $A = D(d, d_1, d_2, d_3)$ corresponds to chopping off the antennas of $\mathcal{T}_A$, if any*. The tropical triangle $\mathcal{T}_{A^{\odot 2}}$ will be called the *soma* of $\mathcal{T}_A$, denoted $\text{soma}(\mathcal{T}_A)$. Then

$$\mathcal{T}_A = \text{soma}(\mathcal{T}_A) \cup \text{ant}(\mathcal{T}_A). \quad (21)$$

Notice the following:

- $\text{soma}(\mathcal{T}_A)$ reduces to one point when $A^{\odot 2} = 0$ (warning: $A^{\odot 2} = 0$ does not imply $A = 0$),
- if $A^{\odot 2} \neq 0$ then $\text{soma}(\mathcal{T}_A)$ is a 2-dimensional convex, compact connected set, in $Z = 0$,
- the antennas of $\mathcal{T}_A$ have integer length d , at most, if $A = D(d, d_1, d_2, d_3)$.

There exist tropical triangles with antennas of arbitrary length. In order to find a canonical normalization for the matrices describing these triangles, we consider

$$E(d, d_1, d_2, d_3, h_1, h_2, h_3) = \begin{bmatrix} 0 & -d - d_2 & -2d - d_3 - h_3 \\ -2d - d_1 - h_1 & 0 & -d - d_3 \\ -d - d_1 & -2d - d_2 - h_2 & 0 \end{bmatrix}. \quad (22)$$

If $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$, then the matrix $E(d, d_1, d_2, d_3, h_1, h_2, h_3)$ is normal. Notice that $E(d, d_1, d_2, d_3, 0, 0, 0) = D(d, d_1, d_2, d_3)$. For pictures in $Z = 0$, we will use

$$E(d, d_1, d_2, d_3, h_1, h_2, h_3)_0 = \begin{bmatrix} d + d_1 & d + h_2 & -2d - d_3 - h_3 \\ -d - h_1 & 2d + d_2 + h_2 & -d - d_3 \\ 0 & 0 & 0 \end{bmatrix}, \quad (23)$$

$$-E(d, d_1, d_2, d_3, h_1, h_2, h_3)^T = \begin{bmatrix} 0 & 2d + d_1 + h_1 & d + d_1 \\ d + d_2 & 0 & 2d + d_2 + h_2 \\ 2d + d_3 + h_3 & d + d_3 & 0 \end{bmatrix}, \quad (24)$$

$$(-E(d, d_1, d_2, d_3, h_1, h_2, h_3)^T)_0 = \begin{bmatrix} -2d - d_3 - h_3 & d + d_1 - d_3 + h_1 & d + d_1 \\ -d + d_2 - d_3 - h_3 & -d - d_3 & 2d + d_2 + h_2 \\ 0 & 0 & 0 \end{bmatrix}, \quad (25)$$

Write $A = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$. Notice that if $h_{j+1} - d_j > 0$, for some j (subscripts modulo 3), the $\mathcal{T}_A$ is not convex in $Z = 0$

Theorem 1 (Canonical normalization). *If A is any 3×3 matrix, then there exist unique $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$, and there exist unique permutation matrices P, Q such that $E(d, d_1, d_2, d_3, h_1, h_2, h_3) = P \odot A \odot Q$. Moreover, $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3).*

Proof. To prove existence, we may assume that A is normal. If $A = A^{\odot 2}$, we take $h_j = 0$, all j , and apply lemma 3. Now assume that $A \neq A^{\odot 2}$. Again, by lemma 3, we can assume that $A^{\odot 2} = D(d, d_1, d_2, d_3)$, for some $d, d_1, d_2, d_3 \geq 0$. Moreover, A is a *tropical square root* of $D(d, d_1, d_2, d_3)$. The geometric meaning of this assertion is that $\mathcal{T}_A$ is obtained from $\mathcal{T}_{A^{\odot 2}}$ by addition of antennas. But for $\mathcal{T}_{A^{\odot 2}}$ to admit adjunction of antennas, d_j must vanish for some j . For each $d_j = 0$ consider any $h_{j+1} > 0$ (subscripts modulo 3). Now, it is easy to verify that the matrix $A = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$ satisfies $A^{\odot 2} = D(d, d_1, d_2, d_3)$.

The uniqueness follows from the geometric meaning of the parameters $d, d_1, d_2, d_3, h_1, h_2, h_3$. $\square$

Corollary 3. *If $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$, and $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3), then $E(d, d_1, d_2, d_3, h_1, h_2, h_3)$ is a tropical square root of $D(d, d_1, d_2, d_3)$.* $\square$

Example 1. *Let us compute the canonical normalization of*

$$A = A_0 = \begin{bmatrix} 0 & 1 & 3 \\ 0 & 3 & 4 \\ 0 & 0 & 0 \end{bmatrix},$$

the example in [7], p. 409. Consider the matrices $U = \text{diag}(-1, 0, 0)$,

$$Q = \begin{bmatrix} -\infty & -\infty & 0 \\ 0 & -\infty & -\infty \\ -\infty & 0 & -\infty \end{bmatrix}$$

and obtain

$$N_0 = U \odot P_{12} \odot A \odot Q = \begin{bmatrix} 2 & 3 & -1 \\ 1 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Then

$$N = N_0 \odot \text{diag}(-2, -3, 0) = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 0 & 0 \\ -2 & -3 & 0 \end{bmatrix}$$

is normal. Then

$$N^{\odot 2} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \\ -2 & -2 & 0 \end{bmatrix}, \quad (N^{\odot 2})_0 = \begin{bmatrix} 2 & 2 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

In figure 14, we find, from left to right, the tropical triangles corresponding to the matrices $A, P_{12} \odot A, T \odot P_{12} \odot A, T \odot P_{12} \odot A \odot Q$ and N (the last three matrices yield the same triangle), all in $Z = 0$. Here $3d = 1, d_1 = d_2 = 0$ and $d_3 = 1$ so that the canonical normalization of $N^{\odot 2}$ is

$$D(1/3, 0, 0, 1) = \begin{bmatrix} 0 & -1/3 & -5/3 \\ -2/3 & 0 & -4/3 \\ -1/3 & -2/3 & 0 \end{bmatrix} = S^{\odot -1} \odot N^{\odot 2} \odot S,$$

with $S = \text{diag}(5/3, 4/3, 0)$. Thus, the canonical normalization of A is

$$S^{\odot -1} \odot N \odot S = E(1/3, 0, 0, 1, 0, 1, 1) = \begin{bmatrix} 0 & -1/3 & -8/3 \\ -2/3 & 0 & -4/3 \\ -1/3 & -5/3 & 0 \end{bmatrix}.$$

We have $h_1 = 0, h_2 = h_3 = 1$. For pictures in $Z = 0$, we consider

$$D(1/3, 0, 0, 1)_0 = \begin{bmatrix} 1/3 & 1/3 & -5/3 \\ -1/3 & 2/3 & -4/3 \\ 0 & 0 & 0 \end{bmatrix}$$

and

$$E(1/3, 0, 0, 1, 0, 1, 1)_0 = \begin{bmatrix} 1/3 & 4/3 & -8/3 \\ -1/3 & 5/3 & -4/3 \\ 0 & 0 & 0 \end{bmatrix}.$$

In figure 15 we see the triangles corresponding to the matrix $N^{\odot 2}$ and its canonical normalization, while in figure 16 we see the triangles of the matrix N and its canonical normalization.

Notice that the matrices $S^{\odot -1}$ and S provide the canonical normalization of $N^{\odot 2}$ and also of N . This holds in general, due to the equality (28) below.




Figure 14: Tropical triangles $\mathcal{T}_A$, $\mathcal{T}_{P_{12} \odot A}$ and $\mathcal{T}_N$.




Figure 15: Tropical triangle corresponding to $N^{\odot 2}$ and to its canonical normalization.




Figure 16: Tropical triangle corresponding to N and to its canonical normalization.

Consider $A = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$, such that $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$ and $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3). Now, a definition of *soma* and *antennas* of $\mathcal{T}_A$ can be given, as in p. 19. The soma of $\mathcal{T}_A$ is $\mathcal{T}_{D(d, d_1, d_2, d_3)}$. The antennas of $\mathcal{T}_A$ have tropical length h_j , if $h_j > 0$.

For any 3×3 real matrix A let $N = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$ be the canonical normalization of A , for some $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$, and $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3). There exist permutation matrices P, Q such that $N = P \odot A \odot Q$. The map f_P is a translation, so that the triangle $\mathcal{T}_A = f_{P \odot -1}(\mathcal{T}_N)$ is just a translated of $\mathcal{T}_N$. Then we define the *soma* and *antennas* of $\mathcal{T}_A$ as follows:

$$\text{soma}(\mathcal{T}_A) = f_{P \odot -1}(\text{soma } \mathcal{T}_N), \quad \text{ant}(\mathcal{T}_A) = f_{P \odot -1}(\text{ant } \mathcal{T}_N), \quad (26)$$

so that equality (21) holds true, for any A .

Remark 1. Sometimes a non-idempotent normal matrix $E(e, e_1, e_2, e_3, h_1, h_2, h_3)$ can be further normalized, to obtain a (simpler) normal matrix $D(d, d_1, d_2, d_3)$, with $d \geq 0, -d \leq d_j$, all j .

Say $h_2 > 0, e_1 = 0, e_2, e_3 \geq 0, h_1 = h_3 = 0$. Then

$$E(e, e_1, e_2, e_3, h_1, h_2, h_3) = \begin{bmatrix} 0 & -e - e_2 & -2e - e_3 \\ -2e & 0 & -e - e_3 \\ -e & -2e - e_2 - h_2 & 0 \end{bmatrix}.$$

Then we take $d = e + \frac{h_2}{3}, d_1 = -h_2, d_2 = e_2, d_3 = e_3$ and $P = \text{diag}(d - e - h_2, -d + e, 0)$ to obtain

$$P \odot E(e, e_1, e_2, e_3, h_1, h_2, h_3) = D(d, d_1, d_2, d_3),$$

but the matrix $D(d, d_1, d_2, d_3)$ will be normal if and only if $d + d_1 = e - \frac{2h_2}{3} \geq 0$.

Example 2. Back to example 1, notice that the matrices

$$D(1, -1, -1, 1) = \begin{bmatrix} 0 & 0 & -3 \\ -1 & 0 & -2 \\ 0 & -1 & 0 \end{bmatrix}$$

and $H = \text{diag}(\frac{1}{3}, \frac{2}{3}, 0)$ satisfy

$$H^{\odot -1} \odot E\left(\frac{1}{3}, 0, 0, 1, 0, 1, 1\right) \odot H = D(1, -1, -1, 1).$$

The following theorem is a simple geometric characterization of normality.

Theorem 2. The 3×3 matrix A is normal if and only if the origin belongs to $\text{soma}(\mathcal{T}_A)$, in $Z = 0$.

Proof. If A is normal then, expression (13) tells us that $\text{col}(A_0, j) \in \overline{C_j^0}$, for $j = 1, 2, 3$. Working in $Z = 0$, let S_j be the tropical segment joining $\text{col}(A_0, j - 1)$ and $\text{col}(A_0, j + 1)$ and let t_j be any classical intersection point of S_j with the tropical line L_0 . Then $\text{span}(t_1, t_2, t_3)$ is contained in L_0 and passes through the origin. Let $\tilde{t}_3 \in S_3$ be a perturbation of t_3 such that $t_1, t_2, \tilde{t}_3$ are not tropical collinear and write $\mathcal{T} = \text{span}(t_1, t_2, \tilde{t}_3)$. Then $\mathcal{T}$ is 2-dimensional, by equality (2) and $\mathcal{T} \subset \mathcal{T}_A$. Moreover, the origin belongs to $\text{span}(t_1, t_2) \subset \mathcal{T}$. Perturbing now t_2 and t_1 in S_2 and S_1 provides the result.

If, in $Z = 0$, the origin belongs to $\text{soma}(\mathcal{T}_A)$, a good tropical triangle, then $\text{col}((A^{\odot 2})_0, j) \in \overline{C_j^0}$, for $j = 1, 2, 3$. Therefore, $\text{col}(A_0, j) \in \overline{C_j^0}$, for $j = 1, 2, 3$, showing normality of A . $\square$

Example 3. Let us look at example 1 again. A normalization of A is $N = P \odot A \odot Q$, where $P = \text{diag}(-2, -3, 0)$,

$$N = \begin{bmatrix} 0 & -1 & -2 \\ 0 & 0 & -3 \\ -1 & 0 & 0 \end{bmatrix}, \quad Q = \begin{bmatrix} -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \\ -1 & -\infty & -\infty \end{bmatrix}.$$

We have

$$N_0 = \begin{bmatrix} 1 & -1 & -2 \\ 1 & 0 & -3 \\ 0 & 0 & 0 \end{bmatrix}, \quad N^{\odot 2} = \begin{bmatrix} 0 & -1 & -2 \\ 0 & 0 & -2 \\ 0 & 0 & 0 \end{bmatrix} = (N^{\odot 2})_0.$$

Another normalization of A is $N = P \odot A \odot Q$, where $P = \text{diag}(-1, -3, 0)$,

$$N = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 0 & -3 \\ -2 & 0 & 0 \end{bmatrix}, \quad Q = \begin{bmatrix} -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \\ -2 & -\infty & -\infty \end{bmatrix};$$

in this case,

$$N_0 = \begin{bmatrix} 2 & 0 & 1 \\ 1 & 0 & -2 \\ 0 & 0 & 0 \end{bmatrix}, \quad N^{\odot 2} = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 0 & -3 \\ -1 & 0 & 0 \end{bmatrix},$$

$$(N^{\odot 2})_0 = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 0 & -2 \\ 0 & 0 & 0 \end{bmatrix}.$$




Figure 17: Tropical triangles $\mathcal{T}_A$ and $\mathcal{T}_N$, for two different normalizations N .

Next, we want to define soma and antennas of a co-spanned tropical triangle. Regarding co-span, we choose to work with *co-normal* matrices, i.e., matrices A having non-negative entries and zero diagonal. We can achieve a canonical co-normalization and then define *soma* and *antennas* of a *co-spanned tropical triangle*, in a similar fashion to theorem 1 and definition in p. 23. Then, what is the relationship between $\mathcal{T}_N$ and $\mathcal{T}^N$, $\text{soma}(\mathcal{T}_N)$ and $\text{soma}(\mathcal{T}^N)$, $\text{ant}(\mathcal{T}_N)$ and $\text{ant}(\mathcal{T}^N)$, for a given canonical normal matrix $N = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$?

- If $h_1 = h_2 = h_3 = 0$, then $N = N^{\odot 2}$ is normal and idempotent. By lemma 2, $\mathcal{T}_N$ is good. Repeating the argument in the proof of corollary 1, we have

$$N^T = (N^{\odot 2})^T = \hat{N}^T,$$

meaning that *the columns of $-N^T$ are precisely the vertices of the sides of $\mathcal{T}_N$* , see figure 12. By the max–min duality, the columns of N are the stable intersection points of the tropical lines N_1, N_2, N_3 . Therefore,

$$\mathcal{T}_N = \text{span}(N) = \text{co-span}(-N^T) = \mathcal{T}^N. \quad (27)$$

- If $h_{j+1} > 0$ and $d_j = 0$ for some j , write $N' = N^{\odot 2}$. By the previous item,

$$\text{soma}(\mathcal{T}_N) = \mathcal{T}_{N'} = \mathcal{T}^{N'} = \text{soma}(\mathcal{T}^N), \quad (28)$$

even if $\mathcal{T}_N \neq \mathcal{T}^N$. Moreover, there is a *bijection between the sets of antennas of $\mathcal{T}_N$ and of $\mathcal{T}^N$* (at most, three antennas each) so that we can talk of *corresponding antennas*. For every antenna a of $\mathcal{T}_N$ and corresponding antenna a' of $\mathcal{T}^N$, there exists a unique cell P_a in the cell decomposition $\mathcal{C}^N$ such that $a' \cup a \subset \overline{P_a}$. Indeed, suppose that $a = \overline{s, q}$ is the antenna of $\mathcal{T}_N$ coming from $h_2 > 0$, $d_1 = 0$, with $s = \text{col}((N^{\odot 2})_0, 2) = [d, 2d + d_2, 0]^T$ and $q = \text{col}(N_0, 2) = [d + h_2, 2d + d_2 + h_2, 0]^T$. Then $a' = \overline{s, r}$ is corresponding antenna in $\mathcal{T}^N$, with $r = \text{col}((-N^T)_0, 3) = [d, 2d + d_2 + h_2, 0]^T$. In $Z = 0$, the cell P_a is determined by

$$d < X, \quad X + d + d_2 < Y < X + d + d_2 + h_2. \quad (29)$$

5 The mapping f_A for a $A = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$

Recall that each antenna a of $\mathcal{T}_A$ gives rise to a cell P_a in the cell decomposition $\mathcal{C}^A$ of $\mathbb{TP}^2$, induced by a matrix A , see p. 12.

Theorem 3. *Let $A = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$, with $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$, and $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3). Then*

1. $\text{soma}(\mathcal{T}_A)$ is the set of fixed points of f_A ,
2. $f_A|_{P_a} = a$, for every antenna a of $\mathcal{T}_A$,
3. $f_A|_P$ is the classical projection onto $\mathcal{T}_A$, in the direction of P , for every unbounded 2-dimensional cell P of the cell decomposition $\mathcal{C}^A$, if $P \neq P_a$, $a \in \text{ant}(\mathcal{T}_A)$.

Proof. First suppose that $h_1 = h_2 = h_3 = 0$. Then $A = D(d, d_1, d_2, d_3)$ is normal idempotent and so, by equalities (9) and (27),

$$\text{soma}(\mathcal{T}_A) = \mathcal{T}_A = \mathcal{T}^A = \overline{B^A}. \quad (30)$$

By lemma 1, the set of fixed points of f_A is $\text{span}(A^{\odot 2}) = \text{span}(A) = \mathcal{T}_A$ and this proves part (1). Since $\mathcal{T}_A$ is a tropical triangle without antennas, then part (2) does not apply. In order to prove part (3), let us work in $Z = 0$. We have A_0 and $(-A^T)_0$ as in equalities (19) and (20). Let P be an unbounded cell of the cell decomposition $\mathcal{C}^A$ parallel, say, to the Y direction and let $p = [x, y, 0]^T$ be a point in P . Then y is a big negative real number or $y = -\infty$, so that $A \odot p$ is a tropical linear combination of the first and third columns of A . In particular, $f_A(p)$ belongs to the (lower part of the) boundary of $\overline{B^A}$ and the point $f_A(p)$ is independent of y . More precisely, if P is determined by the conditions

$$-2d - d_3 < X < d + d_1 - d_3, \quad Y < -d - d_3$$

then $f_A(p) = [x, -d - d_3, 0]^T$. And if P is determined by the conditions

$$d + d_1 - d_3 < X < d + d_1, \quad Y < X - 2d - d_1$$

then $f_A(p) = [x, x - 2d - d_1, 0]^T$. This proves part (3).

Now, suppose that $h_{j+1} > 0$ and $d_j = 0$, for some j , say $j = 2$. The associated antenna a and cell P_a have been described in (29). If $p = [x, y, 0]^T \in \overline{P_a}$ then $f_A(p) = [y - d - d_2, y, x - d]^T = [y - x - d_2, y - x + d, 0]^T \in a$, proving assertion (2). In particular, $f_A(a') = a$, where a' is the antenna of $\mathcal{T}^A$ corresponding to a . $\square$

If $A = D(d, d_1, d_2, d_3)$ for $d, d_1, d_2, d_3 \geq 0$, then $f_A = f_A \circ f_A$, by theorem 3, meaning that f_A is some kind of projection. But, in general, f_A is different from the projector map onto $\mathcal{T}_A$ in p. 10. For instance, consider the matrix $A = D(3, 9, 2, 4)$, i.e.,

$$A = \begin{bmatrix} 0 & -5 & -10 \\ -15 & 0 & -7 \\ -12 & -8 & 0 \end{bmatrix}, \quad A_0 = \begin{bmatrix} 12 & 3 & -10 \\ -3 & 8 & -7 \\ 0 & 0 & 0 \end{bmatrix}, \quad p = \begin{bmatrix} -12 \\ 0 \\ 0 \end{bmatrix}.$$

Then

$$\begin{aligned} \rho(p) &= \begin{bmatrix} -12 \\ -27 \\ -24 \end{bmatrix} \oplus \begin{bmatrix} -12 \\ -7 \\ -15 \end{bmatrix} \oplus \begin{bmatrix} -12 \\ -9 \\ -2 \end{bmatrix} = \begin{bmatrix} -12 \\ -7 \\ -2 \end{bmatrix} = \begin{bmatrix} -10 \\ -5 \\ 0 \end{bmatrix} \\ f_A(p) &= \begin{bmatrix} -5 \\ 0 \\ 0 \end{bmatrix} \neq \rho(p). \end{aligned}$$

Finally, we can describe the mapping f_A , for any 3×3 real matrix A . First, we find the canonical normalization $N = E(d, d_1, d_2, d_3, h_1, h_2, h_3) = P \odot A \odot Q$ to obtain $f_N = f_P \circ f_A \circ f_Q$; then we apply theorem 3, knowing that f_P and f_Q are just changes of coordinates.

What happens to f_A , if some entries of the matrix A become $-\infty$, under the assumption that *at least one entry in each row and in each column is real*? All the

results in the paper still hold true, *except for the non-injectivity and non-surjectivity assertions* in p. 12. Indeed, write $N = E(d, d_1, d_2, d_3, h_1, h_2, h_3) = (n_{ij})$ with $d, d_1, d_2, d_3, h_1, h_2, h_3 \geq 0$ and $h_{j+1} > 0$ implies $d_j = 0$, (subscripts modulo 3). Suppose that $n_{ij} = -\infty$, for some $i \neq j$. Then for some k , the point represented by $\text{col}(N, k)$ lies on a boundary component of $\mathbb{TP}^2$. And if $n_{ij} = -\infty$, for all $i \neq j$, then $N = I$ and $\mathcal{T}_N = \mathbb{TP}^2$. Suppose now that $N = P \odot A \odot Q$ is the canonical normalization of A . Then the following are equivalent:

- $N = I$,
- A is a permutation matrix,
- the columns of A represent $[0, -\infty, -\infty]^T$, $[-\infty, 0, -\infty]^T$ and $[-\infty, -\infty, 0]^T$,
- $\mathcal{T}_A = \mathbb{TP}^2$, i.e., f_A is surjective,
- f_A is injective.

Corollary 4. *A tropical linear mapping on $\mathbb{TP}^2$ transforms tropical collinear points into tropical collinear points.* $\square$

For any real 3×3 matrix A , consider the set of points where f_A is injective,

$$S_A = \{q \in \mathbb{TP}^2 : \exists! p \in \mathbb{TP}^2, f_A(p) = q\}.$$

Working on $\mathbb{R}^n$, the set S_A plays an important role in [5]. If $N = E(d, d_1, d_2, d_3, h_1, h_2, h_3)$, then $S_N = B^N$, by theorem 3 and equality (9). And if $N = P \odot A \odot Q$ is the canonical normalization of A , then $S_A = S_{A \odot Q} = f_{P \odot -1}(S_N)$, meaning that S_A is a translated of S_N .

I would like to thank my students Fernando Barbero and Elisa Lorenzo for their interest and support.

References

- [1] M. Akian, R. Bapat and S. Gaubert, *Max-plus algebra*, chapter 25 in *Handbook of linear algebra*, L. Hobgen (ed.) Chapman and Hall, 2007.
- [2] M. Akian, S. Gaubert and C. Walsh, *Discrete max-plus spectral theory*, in [17], 53–77.
- [3] M. Ansola and M.J. de la Puente, *A note on tropical triangles in the plane*, accepted for publication in Acta Math. Sinica (Engl. ser.), (2008).
- [4] F.L. Baccelli, G. Cohen, G.J. Olsder and J.P. Quadrat, *Synchronization and linearity*, John Wiley; Chichester; New York, 1992.
- [5] P. Butkovič, *Simple image set of $(\max, +)$ linear mappings*, Discrete Appl. Math., **105**, (2000), 73–86.

- [6] B. Carré, *Graphs and networks*, Clarendon Press Oxford, 1979.
- [7] G. Cohen, S. Gaubert and J.P. Quadrat, *Duality and separation theorems in idempotent semimodules*, Linear Algebra Appl., **379**, (2004), 395–422.
- [8] R. Cuninghame–Green, *Minimax algebra*, LNEMS, **166**, Springer, 1970.
- [9] R. Cuninghame–Green, in *Adv. Imag. Electr. Phys.*, **90**, P. Hawkes, (ed.), Academic Press, 1995, 1–121.
- [10] R.A. Cuninghame–Green, P. Butkovič, *Bases in max-algebra*, Linear Algebra Appl. **389**, (2004) 107–120.
- [11] M. Develin, B. Sturmfels, *Tropical convexity*, Doc. Math., **9**, (2004), 1–27; Erratum in Doc. Math. **9** (electronic), 205–206 (2004).
- [12] A. Gathmann, *Tropical algebraic geometry*, Jahresbericht der DMV, **108**, n.1, (2006), 3–32.
- [13] S. Gaubert and Max Plus, *Methods and applications of $(\max, +)$ linear algebra*, in R. Reischuk and M. Morvan, (eds.), STACS’97, **1200** in LNCS, 261–282, Lübeck, March 1997, Springer.
- [14] I. Itenberg, G. Mikhalkin and E. Shustin, *Tropical algebraic geometry*, Birkhäuser, 2007.
- [15] M. Joswig, B. Sturmfels and J. Yu, *Affine buildings and tropical convexity*, Albanian J. Math. **1**, n.4, (2007) 187–211.
- [16] H.W. Kuhn, *The hungarian method for the assignment problem*, Naval Res. Logist., **2**, (1955), 83–97.
- [17] G.L. Litvinov, V.P. Maslov, (eds.) *Idempotent mathematics and mathematical physics*, Proceedings Vienna 2003, American Mathematical Society, Contemp. Math. **377**, (2005).
- [18] G. Merlet, *Semigroup of matrices acting on the max-plus projective space*, submitted to special issue of ILAS 2008, Cancún.
- [19] G. Mikhalkin, *What is a tropical curve?*, Notices AMS, April 2007, 511–513.
- [20] C.H. Papadimitriou and K. Steiglitz, *Combinatorial optimization: algorithms and complexity*, Prentice Hall, 1982, and corrected unabridged republication by Dover, 1998.
- [21] J. Richter–Gebert, B. Sturmfels, T. Theobald, *First steps in tropical geometry*, in [17], 289–317.
- [22] B. Sturmfels, *Solving systems of polynomial equations*, CBMS Regional Conference Series in Math. **97**, AMS, Providence, RI, 2002.

- [23] E. Wagneur, *Moduloids and pseudomodules. Dimension theory*, Discr. Math. **98** (1991) 57–73.
- [24] M. Yoeli, *A note on a generalization of boolean matrix theory*, Amer. Math. Monthly **68**, n.6, (1961) 552–557.